

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 4 : C12N 15/00, A61K 37/02 C07G 17/00, C07H 21/04 C07K 3/02, 3/14, 7/08 C07K 13/00, 15/12		A1	(11) International Publication Number: WO 88/ 03949  (43) International Publication Date: 2 June 1988 (02.06.88)
(21) International Application Number: PCT/AU87/00389 (22) International Filing Date: 19 November 1987 (19.11.87)  (31) Priority Application Numbers: PH 9107 PI 2086 (32) Priority Dates: 21 November 1986 (21.11.86) 22 May 1987 (22.05.87) (33) Priority Country: AU  (71) Applicant (for all designated States except US): HOWARD FLOREY INSTITUTE OF EXPERIMENTAL PHYSIOLOGY AND MEDICINE [AU/AU]; University of Melbourne, Parkville, VIC 3032 (AU). (72) Inventors; and (75) Inventors/Applicants (for US only) : BUTKUS, Aldonna [AU/AU]; 69 Heller Street, West Brunswick, VIC 3055 (AU). COGHLAN, John, Paul [AU/AU]; 4 Walmer Street, Kew, VIC 3101 (AU). ROCHE, Peter, John [AU/AU]; 33 Keith Street, Alphington, VIC 3078 (AU).		(74) Agents: SLATTERY, John, Michael et al.; Davies & Collison, 1 Little Collins Street, Melbourne, VIC 3000 (AU).  (81) Designated States: AT (European patent), AU, BE (European patent), CH (European patent), DE (European patent), FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent), US.  Published With international search report.	
(54) Title: THE CS PROTEIN OF THE CORPUSCLES OF STANNIUS  THE INVENTION RELATES TO THE ISOLATION AND SEQUENCE OF THE CS PROTEIN.  SEQUENCE LISTING (Table with 2 columns: Position, Sequence)  POSITION 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 249			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT Austria	FR France	ML Mali
AU Australia	GA Gabon	MR Mauritania
BB Barbados	GB United Kingdom	MW Malawi
BE Belgium	HU Hungary	NL Netherlands
BG Bulgaria	IT Italy	NO Norway
BJ Benin	JP Japan	RO Romania
BR Brazil	KR Democratic People's Republic of Korea	SD Sudan
CF Central African Republic	KR Republic of Korea	SE Sweden
CG Congo	LI Liechtenstein	SN Senegal
CH Switzerland	LK Sri Lanka	SU Soviet Union
CM Cameroon	LU Luxembourg	TD Chad
DE Germany, Federal Republic of	MC Monaco	TG Togo
DK Denmark	MG Madagascar	US United States of America
FI Finland		

WO 88/03949

PCT/AU87/00389

1

RECEIVED  
CENTRAL FAX CENTER

OCT 14 2004

## THE CS PROTEIN OF THE CORPUSCLES OF STANNIUS

This invention relates to the isolation and characterisation of the CS protein from the Corpuscles of Stannius. The invention is also concerned with the molecular cloning and characterisation of the gene

5 sequence encoding the CS protein.

Note: References referred to hereafter are collected at the end of the description.

In 1839 Stannius described yellow gland-like structures scattered through the kidney of the sturgeon  
10 and noted that these structures also occurred in Teleost fishes as two distinct small bodies and suggested that these organs might represent the adrenal glands of fishes. This homology with the adrenal gland of mammals was accepted until more extensive embryological studies  
15 were undertaken by Huot (1898) who described the differing origins of the Corpuscles of Stannius and the adrenal cortex. However, confusion still persisted. Giacomini in 1908 (a and b) described a mass of tissue in the anterior region of the kidney which on the basis of morphological  
20 characteristics resembled the interrenal system which he designated as the anterior interrenal as distinct from the

SUBSTITUTE SHEET

WO 88/03949

PCT/AU87/00389

2

Stannius corpuscles which he designated as the posterior interrenal. Giacomini also made the observation that the corpuscles were present only in those animals where the parathyroids were absent. Vincent (1898) at about the same time had shown that the removal of the Corpuscles of Stannius was not fatal and therefore assumed they were not essential for life. However Petit (1896) had demonstrated that the removal of one corpuscle resulted in compensatory hypertrophy of the remaining one, thus suggesting that the organ might be an endocrine gland. Following these early observations, the function of the Corpuscles of Stannius has fascinated biologists continuously to 1986 and numerous studies have broached this issue.

The corpuscles from various fishes have been extensively examined morphologically both at the light and electron microscopy level (Vincent, 1898; Oliverseau and Fontaine, 1965; Krishnamurthy and Bern, 1969; Ristow and Piepho, 1963; Wendelaar Bonga and Greven, 1975; Wendelaar Bonga et al. 1980 and Bhattacharyya et al. 1982). There is a consensus view that they contain at least one type of secretory granule. Variations with life cycle, with level of maturation, with age and sex as well as species differences have created a confused picture.

Overall, ablation experiments have led to the view that the Corpuscles of Stannius may have an osmoregulatory role, however their close association with the kidney leave open the possibility of surgical damage to the delicate renal area, which may have compounded the issue of function. There is some evidence that they may have a parathyroid hormone like role in calcium metabolism (Millet et al. 1980 and 1981; Lopez et al. 1984), a prolactin-like action in the gill (Ogasawara and Hirano, 1984) and possibly be the source of a renin-like enzyme capable of angiotensin II production (Chester Jones 1966;

**SUBSTITUTE SHEET**

WO 88/03949

PCT/AU87/00389

3

Ogawa and Sokabe, 1982).

Infusion of extracts of Corpuscles of Stannius have been reported variously as having actions on water, sodium, calcium, phosphate and magnesium metabolism.

- 5 The secretory product of the corpuscles of Stannius has until now remained unresolved.

#### SUMMARY OF THE INVENTION

- The CS protein of the Corpuscles of Stannius has been purified to homogeneity. The partial protein sequence of  
10 the purified CS protein was obtained and used to construct synthetic oligonucleotides which were in turn used to isolate DNA encoding the CS protein of the Corpuscles of Stannius. Characterization of this DNA shows that it encodes the CS protein and a precursor thereof comprising  
15 an N-terminal prohormone segment and a signal peptide sequence which are attached to the CS protein sequence.

- The precursor of the CS protein having an N-terminal  
prohormone segment and signal peptide is hereinafter referred to as the prepro CS protein. The precursor of  
20 the CS protein having an N-terminal prohormone segment is hereinafter referred to as the pro CS protein.

Isolation of the CS protein from the Corpuscles of Stannius will enable further investigations to be carried out on the biological activity of this protein.

- 25 According to one aspect of the present invention, there is provided essentially pure CS protein.

- According to a further aspect of the present invention, there is provided the prepro CS protein, the pro CS protein or the signal or prohormone peptides of the  
30 prepro CS protein. The invention is also directed to sub-units or fragments of the prepro CS protein.

According to a still further aspect of the present invention there is provided a gene encoding the prepro CS protein or sub-units or fragments thereof. Particularly,

#### **SUBSTITUTE SHEET**

WO 88/03949

PCT/AU87/00389

4

there is provided DNA sequences which encode the pro CS protein, the CS protein or the signal or prohormone peptides of the prepro CS protein.

The gene encoding the prepro CS protein or sub-units or fragments thereof may be inserted into appropriate transfer vectors. Suitable vectors include bacterial plasmids, yeast plasmids, phage and other viral vectors.

The gene encoding the prepro CS protein or sub-units or fragments thereof may be inserted into appropriate expression vectors for subsequent expression of the prepro CS protein or sub-units or fragments thereof in eukaryotic or prokaryotic host cells.

Suitable prokaryotic host cells include various E. coli species, Pseudomonas and Serratia marcescens.

Suitable eukaryotic host cells include filamentous fungi, various strains of yeast and mammalian cells (both primary and cell lines).

The isolation of DNA encoding the prepro CS protein and sub-units or fragments thereof will enable large amounts of the corresponding protein products to be produced by recombinant DNA methodology.

According to a yet further aspect of the present invention, there is provided a method for the production of the CS protein, said method comprising the steps of:

- (a) inserting the gene encoding CS protein into an expression vector;
- (b) transfecting a cell with the expression vector of step (a); and
- (c) culturing the cell containing the expression vector to produce the CS protein and subsequently recovering said protein.

As the protein sequence of the prepro CS protein has been elucidated by recombinant DNA techniques, the prepro CS protein and fragments or sub-units thereof may be

**SUBSTITUTE SHEET**

WO 88/03949

PCT/AU87/00389

5

produced using conventional protein synthesis techniques (Barany and Merrifield, 1980).

We have synthesized a peptide fragment of the CS protein precursor corresponding to amino acids 16 to 35 of Figure 3, to investigate the physiological role of this protein. Studies utilising this peptide indicate that the CS protein affects ionic activity in vivo, particularly, in increasing sodium excretion and decreasing plasma potassium concentration. This activity is characteristic of the CS protein, and has been exemplified by, but is in no way restricted to a sheep model. Studies we have carried out in fish indicate this protein is also active in decreasing calcium uptake.

The CS protein and peptide fragments thereof have potential as therapeutic agents in the treatment of cardio-vascular disease, renal disease and electrolyte disorders, particularly, oedema, heart failure and high blood pressure. In addition to the above, the N-terminal peptide fragment of the CS protein precursor corresponding to amino acids 16 to 35 of Figure 3, and other fragments of the CS protein, may be useful tools in determining physiological control mechanisms in the kidney, and the control of blood pressure.

According to another aspect of the present invention, there is provided a therapeutic composition comprising the CS protein or peptide fragments thereof in association with a pharmaceutically acceptable carrier or excipient. Such compositions may be useful in the treatment of cardio-vascular disease, renal disease and electrolyte disorders.

According to a further aspect of the present invention, there is provided a method for the treatment of cardio-vascular disease, renal disease and electrolyte disorders, comprising the administration of a

**SUBSTITUTE SHEET**

WO 88/03949

PCT/AU87/00389

6

therapeutically effective amount of the CS protein or peptide fragments thereof, either alone or in association with a pharmaceutically acceptable carrier or excipient.

The isolation and characterization of DNA encoding the prepro CS protein enables labelled DNA or RNA probes to be prepared. These probes, which may contain all or part of the DNA encoding the prepro CS protein, can be used to identify homologous or similar sequences in higher organisms, including man.

10 DNA probes may be labelled isotopically using for example  $^{32}\text{P}$  or  $^{125}\text{I}$ . DNA probes may also be labelled with biotin or avidin, fluorescent or chemiluminescent reagents, or other appropriate labelling molecules.

RNA probes may be produced from the DNA encoding the  
15 CS protein using, for example, the SP6 vector system (Pharmacia Corporation, Piscataway, N.J.).

It is well established that the genetic code contains redundancies, that is, certain amino acids are coded for by more than one codon. Accordingly, the invention  
20 includes DNA encoding the prepro CS protein or sub-units or fragments thereof where the natural codons are replaced by other codons which code for the same amino acid.

According to a further aspect of the present invention, there is provided a method for the isolation of  
25 the CS protein comprising:

- (a) detergent extraction of the Corpuscles of Stannius;
- (b) electrophoretic separation of the detergent extract in a gel matrix; and
- (c) electroelution of the separated CS protein from step  
30 (b) and recovery of the purified CS protein.

According to a still further aspect of the present invention, there is provided a method for the isolation of a gene encoding the prepro CS protein comprising the steps of:

SUBSTITUTE SHEET
------------------



WO 88/03949

PCT/AU87/00389

7

- (i) preparing a hybridization probe based on the protein sequence of the prepro CS protein;
- (ii) screening a cDNA library prepared from mRNA isolated from the Corpuscles of Stannius with the probe; and
- (iii) identifying and isolating those DNA sequences from the cDNA library of step (ii) which hybridize to the probe.

#### DESCRIPTION OF THE DRAWINGS

FIGURE 1 shows a polyacrylamide gel electrophoresis profile of the isoelectric focusing gel section (pI 5.7 - 6.5) of the CS protein after staining with Coomassie Blue; FIGURE 2 is the sequencing strategy for the CS protein mRNA:

- A. Schematic diagram of the CS protein mRNA. The 5' and 3' untranslated regions are indicated by the solid line. The hatched box represents the signal-peptide sequence, the solid box represents the prosegment, and the open box represents the mature protein.
- B. The arrangement of the four positive CS protein cDNA clones.  $\lambda$ CS.1 was isolated from the RNase H cDNA Library and  $\lambda$ CS.2,  $\lambda$ CS.4 were isolated from the S1-nuclease library.

FIGURE 3 shows the nucleotide sequence and deduced amino acid sequence of the eel CS protein precursor; FIGURE 4 shows a hybridization histochemical analysis of the eel Corpus Stannius-kidney region. 6 $\mu$ m sections of eel CS-kidney region after probing with  $^{32}$ P labelled 75 mer oligonucleotide. The CS region is shown in the lower left region of the autoradiograph.

FIGURE 5 shows the effect of renal arterial infusion into 7 conscious sheep of the N-terminal 25 amino acid fragment of the CS proteins (peptide 'U'). Peptide 'U' was infused at a rate of 50  $\mu$ g/h. Graph (A) is a plot

#### **SUBSTITUTE SHEET**

WO 88/03949

PCT/AU87/00389

8

of urinary sodium excretion ( $\mu\text{mol}/\text{min}$ ) against time (hours). Graph (B) is a plot of plasma potassium against time (hours).

5 Definitions:

The term "CS protein" used herein refers to a polypeptide isolated from the Corpuscles of Stannius, having a molecular weight approximately between 28,000 and 42,000 Daltons as determined by SDS-PAGE and an amino acid  
10 sequence substantially corresponding to amino acids 16-246 of Figure 3.

The amino acid sequence of the CS protein depicted in Figure 3 may be varied by the substitution, addition or deletion of one or more amino acids. Variants which  
15 possess ion-transfer activity, characteristic of the CS protein, are included within the term "CS protein". Such variants may be produced by solid phase peptide synthesis techniques (Barany and Merrifield, 1980). Alternatively,  
20 genetic engineering techniques such as site directed mutogenesis (Botstein and Shortle, 1985), restriction endonuclease digestion, and the ligation of DNA fragments, (Maniatis et al., 1982) may be employed to construct DNA expression vectors which express variants of the CS protein.

25 We have isolated the CS protein from the eel, *Anguilla australis*. It is to be understood that the CS protein may be readily isolated from other animal species following the teaching of the present application. Any such protein having a molecular weight approximately between 28,000 and  
30 42,00 Daltons and an amino acid sequence exhibiting substantial homology with amino acids 16-246 of Figure 3 is to be included within the definition "CS protein".

Substantial homology when used in the above context refers to at least 75% homology with amino acids 16-246 of

**SUBSTITUTE SHEET**

WO 88/03949

PCT/AU87/00389

9

Figure 3.

The term "pro CS protein" used herein, refers to a protein having a sequence corresponding to amino acids 1-246 of Figure 3.

- 5       The term "prepro CS protein" used herein, refers to a protein having a sequence corresponding to amino acids -17-246 of Figure 3.

- 10       "Essentially pure" when used to define the CS protein produced by the present invention refers to the CS protein substantially free of protein or other materials ordinarily associated with the Corpuscles of Stannius, ordinarily greater than or equal to 95% of the total protein being CS protein by weight.

- 15       The term "sub-unit" or "fragment" when used in relation to the CS protein refers to a peptide having an amino acid sequence which is included within the amino acid sequence of the CS protein. Peptides having more than six amino acids are likely to be unique to the CS protein. In order to test whether a peptide is unique to  
20       the CS protein, its amino acid sequence may be compared with amino acid sequences on record in amino acid sequence data banks such as the EMBL Data Base (compiled by the European Molecular Biology Laboratory), the Dayhoff Data Base or the Gene Bank Data Base (compiled by the National  
25       Institutes of Health, U.S.A.). Sub-units or fragments of the CS protein may or may not possess biological activity.

The term "natural codons" refers to those codons which naturally encode amino acids of the prepro CS protein.

- 30       The term "sub-unit" or "fragment", when used in relation to genes encoding the prepro CS protein, refers to a DNA or RNA sequence included in the gene encoding the prepro CS protein. A fragment may comprise single or double stranded DNA or RNA, and is generally in excess of 10 nucleotides. The sequence of any such fragment may be

**SUBSTITUTE SHEET**

WO 88/03949

PCT/AU87/00389

10

compared with sequences recorded in any of the data banks mentioned above to establish whether it is unique to the CS protein.

The following terms have the meanings set forth below:

- 5           DNA    - deoxyribonucleic acid  
          RNA    - ribonucleic acid  
          cDNA   - complementary DNA (enzymically  
                  synthesized from a mRNA sequence  
          mRNA   - messenger RNA  
10          A     - Adenine  
          T     - Thymine  
          G     - Guanine  
          C     - Cytosine  
          U     - Uracil  
15          SDS-PAGE - Sodium dodecyl sulphate poly-  
                  acrylamide gel electrophoresis

EXAMPLES:

EXAMPLE A - Tissue Extraction and Isolation of the CS Protein

20

(i) Tissue Extracts

Corpuscles of Stannius and samples of several other tissues (kidney, atria, ventricle, muscle and liver) were obtained from decapitated eels. These tissues were placed immediately on dry ice and stored under liquid nitrogen until used. A modified Baier technique (1984) was used for homogenation of the tissues. The zwitterionic detergent CHAPS was substituted for Nonidet P40. The tissue homogenates were centrifuged in an Eppendorf (Trade  
25           Mark) centrifuge for 3 minutes and the supernatant used immediately for electrophoresis analysis.

30

(ii) Two dimensional gel electrophoresis techniques

This was performed essentially as originally described by O'Farrell (1975) except that CHAPS was substituted for

**SUBSTITUTE SHEET**

WO 83/03949

PCT/AU87/00389

11

Nonidet P40 as above, and for the electrofocussing gels, the urea solution was deionized just prior to use. After electrofocussing, the gels were equilibrated in O'Farrell's buffer "O" for 10 minutes and then run in the second dimension. These procedures revealed major proteins of the corpuscles of Stannius which are not present in other eel tissues. The portions of the CS electrofocussing gels (pI 5.7 - 6.5) were cut out, equilibrated and run in the second dimension as before. The line of spots containing the CS protein was visualized with 4 M sodium acetate (Higgins and Dehmus, 1979) or Coomassie Blue (CBB) stained (Figure 1).

Once identified the spots were punched out, the gel disc was washed in water and stored at -20°C until electroelution. Electroelution was performed at 5°C according to the technique of Hunkapillar et al (1983) in an electroelution dialysis cell (Caltech).

(iv) Characteristics of the CS Protein

The eluted protein was harvested and precipitated overnight with 9 volumes of cold methanol. Following centrifugation at 5°C, the pellet was washed with a small amount of cold methanol, air dried, dissolved in 0.05% SDS and reprecipitated as above. The air dried pellet was then stored in a dessicator at 5°C until analysed. Electrophoretic analysis has established that the CS protein has a native molecular weight of 42,000, and under reducing conditions a major component of 32,000.

The following results were obtained on analysis of the isolated CS protein:

1. Treatment with N-Glycanase demonstrated that the 32,000 monomer is N-linked glycosylated either at a -Asn-X-Thr or -Asn-X-Ser (where X is any amino acid except Pro) position and that the molecular weight was reduced to 28,000 following removal of the

SUBSTITUTE SHEET
------------------

WO 88/03949

PCT/AU87/00389

12

carbohydrate.

2. Amino acid analysis was performed according to standard procedures on a Beckman System 6300 Amino Acid Analyser.

5 An approximation of the ratio of the amino acid residues in the isolated CS protein (based on an assumed total number of amino acids of 210) gave the following results:

10 Asx (18), Thr (10), Ser (17), Glx (25), Pro (10), Gly (21), Ala (17), Cys (present), Val (15), Met (5), Ile (14), Leu (21), Tyr (10), Phe (13), His (3), Lys (8), Arg (8), indicating a predominance of Glx (Glu or Gln), Gly, Leu, Asx (Asp or Asn), Ser, Ala and the presence of cysteine residues.

- 15 3. Amino acid sequencing

The following partial N-terminal sequence was obtained using an Applied Biosystems gas phase amino acid sequencer:

20 N-Phe-Ser-Ala-Ser-Ser-Pro-Ser-Asp-Val-Ala-Arg-x-Leu-Asn-Gly-Ala-Leu-Gln-Val-Gly-x-Ser-Ala-Phe-Ala-Leu--, where -x- represents cysteine, threonine or possibly tryptophan.

25 **EXAMPLE B - Isolation and Characterisation of a DNA Encoding the prepro CS Protein**

**(i) Synthetic oligonucleotide probes**

30 An oligodeoxyribonucleotide probe was synthesized (using an Applied Biosystems DNA Synthesizer) corresponding to the predicted cDNA sequence of the amino terminal 25 amino acid sequence with the assumption that residues 12 and 21 were Thr and Trp respectively and using preferred codon choices for fish.

**SUBSTITUTE SHEET**

WO 88/03949

PCT/AU87/00389

13

The 75-mer probe sequence was as follows:

5'-GGCGAAGGCGGACCATCCACCTGCAAGGCTCCGTTTCAGTGTCTCTGGC  
CACGTCGGATGGACAGGAGGCGGAGAA-3'

5 The 75-mer probe was 5' end labelled with  $^{32}\text{P}$   
and used to screen the cDNA clone bank constructed from  
mRNA of the Corpuscles of Stannius. This probe was also  
used for hybridization histochemistry studies.

10 (ii) Cloning and Sequence Analysis of cDNA for the  
Isolated CS Protein

Two different cDNA Libraries were constructed  
from eel CS poly(A)<sup>+</sup> RNA. The first library was derived  
from cDNA which was synthesized using standard  
procedures; oligo(dT)-primed AMV reverse transcriptase  
15 reaction followed by DNA polymerase and S<sub>1</sub>-nuclease  
reactions (Maniatis et al, 1982). The second library was  
derived from cDNA synthesized using the RNase H procedure  
of Gubler and Hoffman (1983), which is known to produce  
full-length cDNA transcripts. Both cDNA libraries were  
20 cloned using the bacteriophage vector  $\lambda$ gt 10 (Huynh et  
al, 1985).

Approximately 5,000 clones from each cDNA library  
were screened with the synthetic 75-mer oligonucleotide  
which was 5'-end labelled with [ $\gamma$ - $^{32}\text{P}$ ]ATP.  
25 Hybridization was carried out in 20% formamide, 50 mM  
NaPO<sub>4</sub> pH 6.8, 1mM Na pyrophosphate, 5 x SSC, 5 x  
Denhardtts, 50  $\mu\text{g/ml}$  salmon sperm. DNA at 42°C. Filters  
were washed (4 times) in 0.2 x SSC at 40°C. Three  
hybridization-positive clones were isolated from the  
S<sub>1</sub>-nuclease cDNA library and one hybridization-positive  
30 clone was isolated from the RNase H library. Analysis of  
these clones by restriction endonuclease digests and  
Southern blotting revealed cDNA insert lengths of 0.32  
( $\lambda$ CS.2), 0.8 ( $\lambda$ CS.4) and 1.1 kb ( $\lambda$ CS.3) for the

SUBSTITUTE SHEET

WO 88/03949

PCT/AU87/00389

14

three S<sub>1</sub>-nuclease clones and 2.2 kb for the one RNase H clone ( $\lambda$ CS.1), as shown in Figure 2. Part A of Figure 2 represents a schematic diagram of the CS protein mRNA. The 5' and 3' untranslated regions are indicated by the solid line. The hatched box represents the signal peptide sequence, the solid box represents the prosegment, and the open box represents the mature protein. The nucleotide sequencing strategy and the linear arrangement of these clones is shown in Figure 2.

10 DNA fragments suitable for dideoxy chain termination sequencing (Sanger et al., 1977) were generated by sonication of the appropriate cDNA inserts (Deininger, 1983). These randomly generated fragments were subcloned into M13mp18. The nucleotide sequence of the CS protein cDNA is shown in Figure 3. The sequence of 15 nucleotide residues 269-343 corresponds to the amino-terminal amino acid sequence region determined for the CS protein. It is interesting to note that the original 75-mer probe, designed using preferred codon 20 choices, has 75% identity with the authentic sequence.

On the basis of methods for predicting secretory signal sequences (von Heijne, 1986), it is assumed that the site of cleavage between the signal peptide and the excreted protein is between the alanine and tyrosine 25 residues, indicated by an arrow in Figure 3. This leaves a 15 amino acid pro-sequence extension at the amino terminus of the isolated CS Protein. The pro-sequence precursor is presumably processed by specific cleavage between the arginine and phenylalanine residues (at 30 positions 15 and 16) to give the mature 231 amino acid CS protein shown in square brackets in Figure 3.

The deduced amino acid sequence of the eel CS protein has an Asn-Ser-Thr sequence at amino acids 44-46. The asparagine at position 44 is presumably the

**SUBSTITUTE SHEET**



WO 83/03949

PCT/AU87/00389

15

carbohydrate attachment site. It is interesting to note that there are a total of 15 cysteine residues in the deduced sequence. If all the Cys residues are involved in cysteine bonds, this leaves an extra cysteine which could be free or linked to another peptide chain through a disulfide bond.

EXAMPLE C - Histochemical Localisation of mRNA Encoding the CS Protein

(i) Hybridization Histochemistry

This was performed on the Corpuscles of Stannius and other eel tissue, using the labelled 75-mer oligonucleotide probe as described by Coghlan et al. 1985. These studies revealed that the mRNA coding for this protein was located specifically in the Corpuscles of Stannius and not in any other eel tissues examined. Figure 4 shows specific labelling of the CS and not the kidney with the 75 mer probe.

EXAMPLE D - Synthesis and Biological Activity of a Fragment of the Pro CS Protein

(i) Synthetic Peptide Synthesis

The following peptides, corresponding to amino acids shown in Figure 3, were synthesized by solid-phase synthesis procedures (Barany and Merrifield, 1980) on an Applied Biosystems Model 430A Automated Peptide Synthesizer:

A	-1-15
B	1-15
C	16-35
D	96-109
E	118-151
F	217-246

Crude peptides were purified by gel filtration and preparative high performance liquid chromatography (HPLC)

**SUBSTITUTE SHEET**

WO 88/03949

PCT/AU87/00389

16

under reducing conditions to homogeneity, as assessed by analytical HPLC and amino acid analysis. The cysteines were left unblocked and were tested for biological activity in the reduced monomeric form.

5 (ii) Biological Activity

Sheep were prepared surgically by removing the ovaries and the right kidney, the carotid artery was exteriorized and cannulas placed into the left renal artery and vein. Synthetic peptide was infused directly  
10 into the renal arterial cannula at 50 µg/hr (12 ml/hr) and urine was collected via a Foley bladder catheter. Blood samples were taken from the carotid artery. Urinary sodium and plasma potassium levels were determined by standard procedures. Only peptide C exhibited biological  
15 activity in this bioassay, and the results for this peptide are presented in Figure 5. All statistical analyses were by two-way analyses of variance. Each line in the Figure represents the data for an individual sheep (seven different sheep), and the shaded area shows the  
20 grouped data of one standard error of the mean on either side of the group mean. The results demonstrate that there is a small, but significant, increase in sodium excretion, and also a decrease in plasma potassium concentration in the sheep following infusion of a  
25 synthetic peptide fragment of the pro CS protein.

The above results indicate that the CS protein and fragments thereof, particularly corresponding to peptide C and peptides synthesized from around this region, have potential in the treatment of cardio-vascular  
30 disease, renal disease and electrolyte disorders; particularly oedema, high blood pressure and heart failure. Although these experiments have been carried out in sheep, it is to be understood that the therapeutic affect of these agents is not restricted to sheep.

SUBSTITUTE SHEET

WO 88/03949

PCT/AU87/00389

17

(iii) An Alternative Assay for Biological Activity

The effect of peptide C, corresponding to amino acids 16-35 of Figure 3, on calcium uptake was tested in the assay described by Wagner et al., 1986. This assay measures inhibitory effects upon branchial calcium uptake and involves monitoring the rate of  $^{45}\text{Ca}$  uptake by juvenile rainbow trout.

Results of this assay are shown in Table 1.

TABLE 1

	Dose	(n)	Uptake	P	% Inhibition
	Control (Saline)	7	2.62±0.44		--
	10µg peptide C (4.37nM)	7	0.67±0.14		0.005 (-74%)
	1µg peptide C (0.44nM)	8	1.25±0.19	P	0.01 (-53%)

These results indicate that the CS protein and a selected peptide fragment thereof have an inhibitory effect on calcium uptake.

(iv) Expression of the CS protein in Escherichia Coli

The CS protein cDNA has been cloned into an E. Coli secretion vector, in order to produce large quantities of the CS protein for structural and biological characterization. The CS protein cDNA clone λCS3 was inserted into the EcoRI site of the vector ompA-2, a system developed by Ghrayeb et al. (1984). This secretion vector contains the coding sequence for the ompA signal peptide. This signal peptide guides the cloned gene product across the cytoplasmic membrane into the periplasmic space of the E.Coli cell. The signal peptide is then cleaved from the cloned gene

**SUBSTITUTE SHEET**

WO 88/03949

PCT/AU87/00389

18

product by a specific signal peptidase. Thus, the gene product produced has the authentic amino terminus of the original protein.

5 The coding sequence for the ompA signal peptide was joined onto the coding sequence for the amino terminus of the mature CS protein, i.e. residue 16 in Figure 3, using the site-specific mutagenesis method described by Morinaga et al. (1984). This secretion vector containing the coding sequence for the ompA signal  
10 peptice/CS protein was designated pCS-2. Gel electrophoresis of proteins from E. Coli cells containing pCS-2, and Western blot analysis using a CS protein antisera, indicate that the cells are producing a gene product which binds the CS protein antibody.

15

**SUBSTITUTE SHEET**

WO 88/03949

PCT/AU87/00389

19

REFERENCES

- 5 Baier, L.J., Hanash, S.M. and Erickson, R.P. (1984)  
Mice homozygous for chromosomal deletions at the  
albino locus region lack specific polypeptides in  
two-dimensional gels.  
Proc. Natl. Acad. Sci. USA 81: 2132-2136
- 10 Barany, G. and Merrifield, R.B., (1980) Solid-Phase  
Peptide Synthesis, In "The Peptides", Vol. 2 (E.  
Gross & J. Meienhofer eds.) Academic Press, New  
York: 1-284
- 15 Bhattacharyya, T.K., Buttler, D.G., Youson, J.H. (1982)  
Ultrastructure of the corpuscles of Stannius in the  
garpike (*Lepisosteus platyrhynchus*).  
Gen. Comp. Endocrinol. 46: 29-41
- 20 Botstein D. and Shortle D., (1985) Strategies and  
Applications of In Vitro Mutagenesis, Science, 229:  
1193-1201
- 25 Chester Jones, I., Henderson, I.W., Chan, D.K.O.,  
Rankin, J.C., Mosley, W., Brown, J.J., Lever, A.F.,  
Robertson, J.I.S. and Tree, M. (1956)  
Pressor activity in extracts of the corpuscles of  
Stannius from the European eel (*Anguilla anguilla*).  
J. Endocrin. 34: 393-408
- 30 Coghlan, J.P., Aldred, P., Haralambidis, J., Niall,  
H.D., Penschow, J.D. and Tregear, G.W. (1985)  
Review of hybridization histochemistry.  
Analytical Biochemistry 149: 1-28
- 35 Deininger, P.L. (1983)  
Random subcloning of sonicated DNA: application to  
shotgun DNA sequence analysis.  
Anal. Biochem., 122: 216-223
- 40 Ghrayeb, J., Kimura, H., Takahara, M., Hsiung, H.,  
Masui, Y. and Inouye, M. (1984) Secretion cloning  
vectoris in *Escherichia coli*, The Embo Journal, 3:  
2437-2442
- 45 Giacomini, E. (1908a)  
Sulla disposizione del sistema interrenale e del  
sistema feocromo nella cieche, e nei leptocephali  
Rendic. Accad. sc. Ist. Bologna, n.s., 12: 172
- 50 Giacomini, E. (1908b)

**SUBSTITUTE SHEET**

WO 88/03949

PCT/AU87/00389

20

If sistema interrenale e il sistema cromaffine  
(sistema feocromo) nelle Anguille adulte, nell eiecha,  
e nei leptocefali.

Mem. Accad. Sc. Ist. Bologna, ser 6, T.5, 407

Abstract in Monit. Zool. Ital. 20: 92

Gubler, U. and Hoffman, B.J. (1983)

A simple and very efficient method for generating cDNA  
libraries.

Gene 25: 263-269

Higgins, R.C. and Dahmus, M.E. (1979)

Rapid visualization of protein bands in preparative  
SDS-polyacrylamide gels.

Anal. Biochem. 93: 257-260

Huot, E. (1898)

Preliminaire sur l'origine des capsules surrenales des  
poissons lophobranches.

Compt. Rend. Acad. d. sc., T.126,49.

Hunkapillar, M.W., Lujan, E., Ostrander, F. and Hood,  
L.E. (1983)

Isolation of microgram quantities of proteins from  
polyacrylamide gels for amino acid sequence analysis.

Methods in Enzymology 91: 227-236

Huynh, T.V., Young, R.A. and Davis R.W. (1985)

Constructing and screening cDNA libraries in  $\lambda$ gt10  
and  $\lambda$ gt 11.

In: DNA Cloning Volume I : a practical approach  
(David Glover, ed.) IRL Press, Oxford, p 49-78

Krishnamurthy V.G. and Bern, H.A. (1969)

Conductive Histologic Study of the Corpuscles of  
Stannius and the Juxtaglomerular Cells of Teleost  
Fishes.

Gen. Comp. Endocrinol. 13: 313-335

Lopez, E., Tisserand-Jochem, E-M., Eyquem, A., Milet,  
C., Hillyard, C., Lallier, E., Vidal, B. and  
MacIntyre, I. (1984)

Immunocytochemical detection in eel corpuscles of  
stannius of a mammalian parathyroid-like hormone.

Gen. & Comp. Endocrinol. 53: 28-36

Maniatis, T., Fritsch, E.F. and Sambrook, J. (1982)

Molecular Cloning : A laboratory manual. Cold Spring  
Harbor Laboratory, ColdSpring Harbor, New York

**SUBSTITUTE SHEET**

WO 88/03949

PCT/AU87/00389

21

- 5 Milet, C., Hillyard, C.J., Martelly, E., Girgis, S.,  
MacIntyre, I. and Lopez, E. (1980)  
Similitudes structurales entre l'hormone  
hypocalcemiante des corpuscules de stannius (PCS) par  
l'Anguille (*anguilla anguilla* L.) et l'hormone  
parathroidienne mammalienne.  
Comptes Rendus Hebd. Seances (Acad. sci. Ser. D. Sci.  
Nat.) 291: (12) 977-980
- 10 Morinaga, &., Franceschini, T., Inouye, S. and Inouye,  
M. (1984), Improvement of oligonucleotide-directed  
site-specific mutagenesis using double-stranded  
plasmid DNA. *Biotechnology*, 2: 636-639
- 15 O'Farrel, P.H. (1975)  
High resolution of two-dimensional electrophoresis of  
proteins.  
*J. Biol. Chem.* 250: (10) 4007-4021
- 20 Ogasawara, T. and Hirano, T. (1984)  
Effects of prolactin and environmental calcium on  
osmotic water permeability of the gills of the eel  
*Anguilla japonica*.  
*Gen. & Comp. Endocrinol.* 53: 315-324
- 25 Ogawa, J. (1968)  
Osmotic and ionic regulation in goldfish following  
removal of the corpuscles of Stannius or the pituitary  
gland.  
30 *Can. J. Zool.* 46: (4) 669-676
- Ogawa, M. and Sokabe, H. (1982)  
Hypocalcemic effect of homologous angiotensin-like  
substances produced by the renin-like enzyme in the  
35 corpuscles of Stannius in the Japanese eel *Anguilla*  
*japonica*.  
*Gen. & Comp. Endocrinol.* 47: 36-41
- 40 Olievereau, M. and Fontaine, M. (1965)  
Effect de l'hypophysectomie sur les corpuscles de  
Stannius de l'Anguille  
*Comp. Rend. Acad. Sci.* 261: 2003-2008
- 45 Petit, A. (1896)  
Recherches sur les capsules surrenales.  
*J. de l'anat. et physiol. T.* 32, 391
- 50 Ristow, H. and Piepho, H. (1963)  
Uber die Bildung der Sekretgranula in den  
Stanniusschen Korperchen des FluBaales.

SUBSTITUTE SHEET

WO 88/03949

PCT/AU87/00389

22

Naturwissenschaften 50: 382-383

5 Sanger, F., Nicklen, S. and Coulson, A.R. (1977)  
DNA sequencing with chain terminating inhibitors.  
Proc. Natl. Acad. Sci. USA 74: 5463-5467

10 Stannius, H. (1839)  
Die Nebennierenbeiknochenfischen.  
Arch. Anat. Physiol. 2: 233-271

15 Von Heijne, G. (1986)  
A new method for predicting signal sequence cleavage  
sites.  
Nucleic Acids Research, 14: 4683-4690

20 Vincent, S. (1898)  
The effects of extirpation of the suprarenal bodies of  
the eel (*Anguilla anguilla*).  
Proc. Roy. Soc. London, 52: 354-356

25 Wagner, G.F., Hampong, M., Park, C.M. and Copp, D.H.  
(1986) Purification, Characterization, and Bioassay  
of Teleocalcin, a Glycoprotein from Salmon Corpuscles  
of Stannius. General and Comparative Endocrinology  
53: 481-491

30 Wendelaar-Bonga, S.E. and Greven, J.A.A.  
A Second Cell Type in Stannius Bodies of Two  
Euryhaline Teleost Species.  
Cell. Tiss. Res., 159: 287-290

35 Wendelaar-Bonga, S.E., van der Meij, J.C.A. and Pang,  
P.K.T. (1980)  
Evidence for two secretory cell types in the Stannius  
bodies of the Teleosts *Fundulus heteroclitus* and  
*Carassius auratus*.  
Cell & Tissue Research 212: 295-306

40

SUBSTITUTE SHEET



WO 88/03949

PCT/AU87/00389

23

## CLAIMS:

1. Essentially pure CS protein.
2. Essentially pure pro CS protein.
3. Essentially pure prepro CS protein.
4. The signal or prohormone peptide of the prepro CS protein.
5. A peptide fragment of the pre pro CS protein.
6. A fragment as claimed in claim 5, which possess ion transfer activity characteristic of the CS protein.
7. A fragment as claimed in claim 6 having a sequence corresponding to amino acids 16 to 35 of Figure 3.
8. A therapeutic composition comprising the CS protein or peptide fragments thereof in association with a pharmaceutically acceptable carrier or excipient.
9. A method for the treatment of cardio-vascular disease, renal disease or electrolyte disorders, comprising the administration of a therapeutically effective amount of the CS protein or peptide fragments thereof either alone or in association with a pharmaceutically acceptable carrier or excipient.
10. A method for the isolation of the CS protein comprising:
  - (a) detergent extraction of the Corpuscles of Stannius;

**SUBSTITUTE SHEET**

WO 88/03949

PCT/AU87/00389

24

(b) electrophoretic separation of the detergent extract; and

(c) electroelution and recovery of the separated CS protein from step (b).

11. A gene encoding the prepro CS protein.

12. A sub-unit of the gene claimed in claim 11, which encodes the pro CS protein, the CS Protein or the signal or prohormone peptides of the prepro CS protein.

13. A fragment of the gene claimed in claim 11, excluding those sub-units claimed in claim 12, which is unique to the gene encoding the prepro CS protein.

14. A transfer vector containing a gene, sub-unit or fragment as claimed in any one of claims 11 to 13.

15. A method for the isolation of a gene encoding the prepro CS protein comprising the steps of:

(i) preparing a hybridization probe based on the protein sequence of the prepro CS protein;

(ii) screening a cDNA Library prepared from mRNA extracted from the Corpuscles of Stannius with the probe; and

(iii) identifying and isolating those DNA sequences from the cDNA Library of step (ii) which hybridize to the probe.

16. A method for the production of the CS protein, said method comprising the steps of:

(a) inserting the gene encoding CS protein into an expression vector;

(b) transfecting a cell with the expression vector of

**SUBSTITUTE SHEET**

WO 88/03949

PCT/AU87/00389

25

- step (a); and
- (c) culturing the cell containing the expression vector to produce the CS protein and subsequently recovering said protein.

17. A method as claimed in claim 16, wherein the expression vector is a viral vector.

18. A method as claimed in claim 16, wherein the vector is ompA-2 and the transfected cell is E. Coli.

19. The CS protein when produced by the methods of any one of claims 16 to 18.

20. The CS protein, methods for its preparation and pharmaceutical compositions containing it substantially as hereinbefore described.

21. A gene encoding the CS protein and methods for its production substantially as hereinbefore described.

SUBSTITUTE SHEET

WO 88/03949

PCT/AU87/00389

1/9



*Fig. 1.*

**SUBSTITUTE SHEET**

WO 88/03949

PCT/AU87/00389

2/9




FIG.2.

SUBSTITUTE SHEET

WO 88/03949

PCT/AU87/00389

THE NUCLEOTIDE SEQUENCE AND DEDUCED AMINO ACID SEQUENCE OF THE ed<sub>1</sub> CS PROTEIN PRECURSOR.

5'-GATAACACCGGTAAACAATCGTCACGTGAGGTCTAAACTACAACTTTCTGAACACAGACAAGAAACCGCCTTGCGCCAT 80  
10 20 30 40 50 60 70  
TTGTTGTCACATCCAGAAACCTTAAGTATCTTGTCTTAAATG 120  
90 100 110 120  
Thr Leu Val Leu Val Thr Ala Ala Tyr Glu Gln Asp Glu Ser Glu Pro Leu Ser Pro Arg  
ACG CTT GTG CTG GTA ACT GCT GCC TAC TAC GAG CAG GAT GAG AGC GAG CCC TTA TCT CCA AGG 208  
15 20 25 30 35 40 45 50 55 60 65 70  
Thr Ala Arg Phe Ser Ala Ser Ser Pro Ser Asp Val Ala Arg Cys Leu Asn Thr Asp Gly Ala Leu  
ACA GCG CGC TTC TCC GCC AGC AGC CCA TCT GAT GTT GCA CGC TGT CTG AAC GGG GCC CTG 268  
Gln Val Gly Cys Ser Ala Phe Ala Cys Leu Asp Asn Thr Cys Asn Thr Asp Gly Met  
CAG GTG GGC TGC AGT GCA TTT GCC TGT TGT GAC GAC CTT CTT GGT GAC ACC AAC ACC 318  
35 40 45 50 55 60 65 70  
His Glu Ile Cys Arg Ser Phe Leu His Gly Ala Ala Lys Phe Asp Thr Gln Gly Lys Thr  
CAT GAA ATC TGC AGG TCC TTC CTC CAC CAC GGT GCT GCC AAA TTT GAC ACA CAG GGC AAG ACT 388  
100 110 120 130 140 150 160 170 180 190 200 210 220 230 240 250 260 270 280 290 300 310 320 330 340 350 360 370 380 390 400

FIG.3Aa.

SUBSTITUTE SHEET

WO 88/03949

PCT/AU87/00389

4/9

75	Phe	Val	Lys	Glu	Ser	Leu	Lys	Ile	Ala	Asn	Gly	Ile	Thr	Ser	Lys	Val	Phe	Thr	Ala
	TTT	GTG	AAG	GAG	AGC	CTG	AAG	TGC	ATA	AAT	GGC	ATC	ACC	TCC	AAA	GTG	TTC	CTT	ACC
	398		408			418					428				438		90		448
95	Ile	Arg	Arg	Cys	Ser	Ser	Phe	Gln	Lys	Met	Ile	Glu	Val	Gln	Glu	Glu	Cys	Tyr	Ser
	ATC	CGC	CGC	TGC	TCA	TCC	TTC	CAG	AAG	ATG	ATC	GAG	GTT	CAG	GAG	GAG	TGC	TAT	AGC
	458		468			478					488				498		115		508
115	Lys	Leu	Asp	Leu	Cys	Ser	Val	Ala	Gln	Ser	Asn	Glu	Ala	Met	Gly	Glu	Val	Ala	Gln
	AAA	CTA	GAC	CTC	TGC	TCT	GTT	GCC	CAG	AGC	AAC	GAG	GCC	ATG	GAG	GAG	GTG	GCC	CAG
	518		528			538						548			558		130		568
135	Val	Pro	Ser	Gln	Phe	Pro	Asn	Arg	Tyr	Tyr	Ser	Leu	Leu	Gln	Ser	Leu	Leu	Thr	Cys
	GTG	CCC	AGC	CAG	TTT	CCC	AAC	AGG	TAC	TAC	AGC	CTG	CTG	CAG	AGT	CTT	CTG	ACG	TGT
	578		588			598						608			618		150		628
155	Asp	Glu	Asp	Thr	Val	Glu	Gln	Val	Arg	Ala	Gly	Val	Ser	Arg	Leu	Glu	Pro	Glu	Met
	GAT	GAG	GAC	ACC	GTG	GAG	CAG	GTG	AGG	GCC	GGG	TTC	TCC	CGC	CTG	GAG	CCA	GAG	ATG
	638		648			658						668			678		170		688

FIG. 3Ab.

SUBSTITUTE SHEET

WO 88/03949

PCT/AU87/00389

5/9

Gly Val GTG	175	Gln Phe	180	Cys Thr	185	Pro Pro Ser Ala Ala	190	Gly Gly Thr	748
GGG		CTC TTC	708	TGC		CCG AGC GCG GCG	738	GGT GGT ACT	
Gly Pro Val	195	Gly Ala Gly	200	Cys Thr	205	Pro Gly Pro Cys	210	Ser Arg Ser	808
GGC CCA		GTA GGG GCA GGA	758	TGC		TGG GGC CCC CCA	788	TGT TCA AGA	
Ser Pro Thr	215	Cys Ala Pro	220	Thr Thr	225	Leu Leu Arg Asn Ala	230	Arg Pro Ala	868
AGC CCA		ACC TGC GCT CCC	818	ACC		TGT TTTG CTA AGA AAC	848	GCT CCA GCT	
Pro Asn Tyr	235	His Pro Pro	240	Met Asp	245	Cys Pro	908	Pro CCA	928
CCT AAT		TAC CAC CCA CCA	878	GAT		TGT CCT	918	TAA	
GTAGAGAACAAATTGTGCTATTTCCTCTTATGTACACAGGGGTAATAAGCTTCCAACTAGGTTTTTAAACAAACAAAC	942	952	962	972	982	992	1002	1012	

FIG. 3AC.

SUBSTITUTE SHEET



WO 88/03949

PCT/AU87/80389

## CONTINUATION OF 3' UNTRANSLATED REGION OF THE eel CS NUCLEOTIDE SEQUENCE.

CTTCAGTAAACACATGTGGTCTACTAGACGACAGTATTTGATGGTGGACCGTGTATGTAATGTAAGTACTGGACACT  
1022 1032 1042 1052 1062 1072 1082 1092

CAGCTTGTAAATGTGAAGCTTTGTGTTTCTTCAATGGTACAGTATATTCATTTCTTTTATGTAATGCCCTTGTATATCAT  
1102 1112 1122 1132 1142 1152 1162 1172

CAAGTTTGTAAAGAGAATATCAGGCTGTCATGGGTTTATTTTGTGAAGGTACATTAATATATATATATATATATATATATTT  
1182 1192 1202 1212 1222 1232 1242 1252

GAGTTTAAAAATATTGTCCTTGTACAAATCAAGGATTTTACCGAGCATAGCCAAATCACAATCTTTTCAATCTTTTTCCTCAA  
1262 1272 1282 1292 1302 1312 1322 1332

ACCTCAAAGCAGATATTGAACAACATGCTCTTTGGAAATGAGGATAATCAGGATCCCTGTTCTGATAATGCAACCAATAAA  
1342 1352 1362 1372 1382 1392 1402 1492

GAGACCTAACAGGACGATTCTGTAGTCTTTGTTAATAATTTTATTTGATATTTAACTTCTTAATAATAATAATAATAATAATA  
1422 1432 1442 1452 1462 1472 1482 1492

TAATGTTTTTATTTGGTTGAGTTGCCATGGTTACTATTTGTAAGTACCCAGGAGTTGAATGCGCAGTTCCTGTTTCAGCCCT  
1502 1512 1522 1532 1542 1552 1562 1572

CGTGGTGAATCTTGAAGCCCACTCAGAGGCGAGTTAATTCATTAATGAACAGGACATGTGCTTCGATCCCATGCTGGTCCGAT  
1582 1592 1602 1612 1622 1632 1642 1652

SUBSTITUTE SHEET

FIG. 3Ba.

WO 88/03949

PCT/AU87/00389

7/9

GAGGGTCATTACAGCAAAACATATTGGCACAAATTATCTGCCGGTATTTTCCAGGGCCCTTATATTAGACAGCTCAGCAACT  
1662 1672 1682 1692 1702 1712 1722 1732

TGGCAAGGTTTGGAAACCTTTGTAAAGACAGAAAGGTCACAAACATGTCAATAGTGTACCTCGTGTGCCATAACACAGA  
1742 1752 1762 1772 1782 1792 1802 1812

TTGATTAGCTCTATACATTTTGTAAAGTACAAACTTTCTGGTGGTTTGCCTGTGCAAAATAC TACTAATAACAGGTCAT  
1822 1832 1842 1852 1862 1872 1882 1892

AATAGCAAGAGAGACTCTTCTACACCCAGCCCTGATTGGCTAAGGAAACAC TGTGCAGATTGTGATAGGATGCTTCATAC  
1902 1912 1922 1932 1942 1952 1962 1972

TGTTGCCCTGATATGCC TGATTTTAAGGCCCTTAGCCAAACTCAGCAGTTGATTTCTTTGTTTGC TTTTGAGCAAAAGTTAAG  
1982 1992 2002 2012 2022 2032 2042 2052

GCCTAAAGCACCTTGAAGTCAGATTCCAGAATTAACTTCAATGATTTGCTTCAAGGACTATTGTCATGCTTTTGGGGGCA  
2062 2072 2082 2092 2102 2112 2122 2132

AAACCTTGGT TGAGATCGAGATAATGTCCACACATTAAAGAAAATGCCAAAAAAACCGG-3'  
2142 2152 2162 2172 2182


Fig. 3Bb.

WO 88/03949

PCT/AU87/00389

8/9

Fig. 4.



SUBSTITUTE SHEET

WO 88/03949

PCT/AU87/00389

9/9




FIG. 5 (A).




FIG. 5 (B).

SUBSTITUTE SHEET



International Application No. PCT/AU 87/00389

## FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

Volume 240, issued 1986 (Alan R. Liss, Inc., New York, USA) see pages 363 to 367

- A Chemical Abstracts, Volume 103, no. 17, issued 1985, October 28, (Columbus, Ohio, USA), Takagi, Y. et al. "Effects of the removal of corpuscles of Stannius on the transport of calcium across the intestine of rainbow trout", see page 463, column 2, the abstract no. 138988e, Zool. Sci., 1985, 2 (4), 523 to 530.

(continued)

V. ☒ OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

- ☒ Claim numbers 9 because they relate to subject matter not required to be searched by this Authority, namely:

it involves a method of treatment of the human or animal body (Rule 39).

- ☐ Claim numbers ..... because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

- ☐ Claim numbers ..... because they are dependent claims and are not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☐ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING

This International Searching Authority found multiple inventions in this international application as follows:

- ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.
- ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:
- ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims if it is covered by claim number(s):
- ☐ As all searchable claims could be searched without effect paying an additional fee, the International Searching Authority did not invite payment of any additional fee.

## Remarks on Protest

- ☐ The additional search fees were accompanied by applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

Form PCT/ISA/219 (Supplemental sheet (2)) (January 1993)

International Application No. PCT/AU 87/00389

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
A	Chemical Abstracts, Volume 99, no. 19, issued 1983, November 7, (Columbus, Ohio, USA). Fenwick, J.C. "Some evidence concerning the nature of the hypocalcemic factor in the Stannius corpuscles", see page 374, column 2, the abstract no. 155575j, Comp. Endocrinol. Calcium Regul., Proc. Satell. Symp. 1981 (Pub. 1982), 167 to 72.	
P,A	Harvey, S. et al. "Parathyroid Hormone-like Immunoreactivity in Fish Plasma and Tissues", General and Comparative Endocrinology, Volume 68, issued 1987, (Academic Press, Inc., New York, USA), see pages 136 to 146.	

Form PCT ISA 210 (extra sheet) (January 1983)